厦门大学高等代数教案 网站 IP 地址: 59.77.1.116; 域名: gdjpkc.xmu.edu.cn

§1.5 行列式的计算

教学目的与要求 利用行列式性质,掌握计算行列式的值的典型方法,掌握用数学归纳法求行列式值.

例 1

$$|A| = \begin{vmatrix} 3 & 0 & 15 & 12 \\ 1 & 2 & 5 & 6 \\ 2 & 0 & -3 & 0 \\ 5 & 0 & 1 & 2 \end{vmatrix}$$

解

$$|A| = 2 \begin{vmatrix} 3 & 15 & 12 \\ 2 & -3 & 0 \\ 5 & 1 & 2 \end{vmatrix} = 6 \begin{vmatrix} 1 & 5 & 4 \\ 2 & -3 & 0 \\ 5 & 1 & 2 \end{vmatrix} = 6 \begin{vmatrix} -9 & 3 & 0 \\ 2 & -3 & 0 \\ 5 & 1 & 2 \end{vmatrix} = 12 \begin{vmatrix} -7 & 0 \\ 2 & -3 \end{vmatrix} = 3 \times 21 = 252$$

例 2 设 $a_i \neq 0$, $0 \leq i \leq n$, 求

$$|A| = \begin{vmatrix} a_0 & b_1 & \cdots & b_n \\ c_1 & a_1 & & & \\ \vdots & & \ddots & & \\ c_n & & & a_n \end{vmatrix}.$$

解(法1)消去第1列

$$|A| = \begin{vmatrix} a_0 - \sum_{i=1}^n \frac{b_i c_i}{a_i} & b_1 & \cdots & b_n \\ & a_1 & & \\ & & \ddots & \\ & & & a_n \end{vmatrix} = (\prod_{i=1}^n a_i)(a_0 - \sum_{i=1}^n \frac{b_i c_i}{a_i})$$

(法2)按第1列展开

$$|A| = \prod_{i=0}^{n} a_i - \sum_{i=1}^{n} b_i c_i \prod_{j=1, j \neq i}^{n} a_j$$

例 3 求

$$F_n = \begin{vmatrix} \lambda & 0 & 0 & \cdots & 0 & a_n \\ -1 & \lambda & 0 & \cdots & 0 & a_{n-1} \\ 0 & -1 & \lambda & \cdots & 0 & a_{n-2} \\ \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda & a_2 \\ 0 & 0 & 0 & \cdots & -1 & \lambda + a_1 \end{vmatrix}$$

解 (法 1) 按第 1 列展开. $F_n = \lambda F_{n-1} + (-1)^{1+n} (-1)^{n-1} a_n = \lambda F_{n-1} + a_n$. 从而 $F_n = \lambda^n + a_1 \lambda^{n-1} + \dots + a_n$.

(法2) 按第 n 列展开

$$F_n = (-1)^{1+n} a_n \begin{vmatrix} -1 & \lambda & 0 & \cdots & 0 \\ 0 & -1 & \lambda & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & \lambda \\ 0 & 0 & 0 & \cdots & -1 \end{vmatrix} + (-1)^{2+n} a_{n-1} \begin{vmatrix} \lambda & 0 & 0 & \cdots & 0 \\ 0 & -1 & \lambda & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & \lambda \\ 0 & 0 & 0 & \cdots & -1 \end{vmatrix}$$

$$+\cdots+(\lambda+a_1)\begin{vmatrix}\lambda&0&\cdots&0\\-1&\lambda&\cdots&0\\\cdots&\cdots&\cdots&\cdots\\0&0&\cdots&\lambda\\0&0&\cdots&-1\end{vmatrix}$$

$$= (-1)^{1+n}(-1)^{n-1}a_n + (-1)^{2+n}(-1)^{n-2}\lambda a_{n-1} + \dots + (\lambda + a_1)\lambda^{n-1}$$

= $a_n + a_{n-1}\lambda + \dots + a_1\lambda^{n-1} + \lambda^n$

(法 3)

$$F_{n} = \begin{vmatrix} \lambda & 0 & 0 & \cdots & 0 & 0 & a_{n} \\ -1 & \lambda & 0 & \cdots & 0 & 0 & a_{n-1} \\ 0 & -1 & \lambda & \cdots & 0 & 0 & a_{n-2} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & -1 & 0 & a_{2} + (\lambda + a_{1})\lambda \\ 0 & 0 & 0 & \cdots & 0 & -1 & \lambda + a_{1} \end{vmatrix} = \cdots$$

$$= \begin{vmatrix} 0 & a_{n} + a_{n-1}\lambda + \cdots + a_{1}\lambda^{n-1} + \lambda^{n} \\ -1 & 0 & \vdots \\ & \ddots & 0 & a_{2} + a_{1}\lambda + \lambda^{2} \\ & -1 & \lambda + a_{1} \end{vmatrix}$$

$$= (-1)^{n-1}(-1)^{n+1}(a_{n} + a_{n-1}\lambda + \cdots + a_{1}\lambda^{n-1} + \lambda^{n})$$

$$= a_{n} + a_{n-1}\lambda + \cdots + a_{1}\lambda^{n-1} + \lambda^{n}$$

例 4

$$|A| = \begin{vmatrix} x & a & a & \cdots & a \\ a & x & a & \cdots & a \\ a & a & x & \cdots & a \\ \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ a & a & a & \cdots & x \end{vmatrix}$$
 $(n > 1)$

解 (法 1) 第 2 行, 第 3 行, \cdots , 第 n 行加到第 1 行上

$$|A| = \begin{vmatrix} x + (n-1)a & x + (n-1)a & x + (n-1)a & \cdots & x + (n-1)a \\ a & x & a & \cdots & a \\ a & a & x & \cdots & a \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a & a & a & \cdots & x \end{vmatrix}$$

$$= [x + (n-1)a] \begin{vmatrix} 1 & 1 & 1 & \cdots & 1 \\ a & x & a & \cdots & a \\ a & a & x & \cdots & a \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ a & a & a & \cdots & x \end{vmatrix}$$

$$= [x + (n-1)a] \begin{vmatrix} 1 & 1 & 1 & \cdots & 1 \\ 0 & x - a & 0 & \cdots & 0 \\ 0 & 0 & x - a & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & x - a \end{vmatrix}$$

$$= [x + (n-1)a](x - a)^{n-1}$$

(法2)

$$|A| = \begin{vmatrix} x & a & a & \cdots & a \\ a - x & x - a & 0 & \cdots & 0 \\ a - x & 0 & x - a & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ a - x & 0 & 0 & \cdots & x - a \end{vmatrix}$$

$$= \begin{vmatrix} x + (n-1)a & a & a & \cdots & a \\ 0 & x - a & 0 & \cdots & 0 \\ 0 & 0 & x - a & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & x - a \end{vmatrix}$$

$$= [x + (n-1)a](x-a)^{n-1}$$

(法 3) 加边法.

$$|A| = \begin{vmatrix} 1 & a & a & a & \cdots & a \\ 0 & x & a & a & \cdots & a \\ 0 & a & x & a & \cdots & a \\ 0 & a & a & x & \cdots & a \\ \vdots & \vdots & \ddots & \ddots & \vdots & \ddots & \vdots \\ 0 & a & a & a & \cdots & x \end{vmatrix}_{n+1}$$

$$= \begin{vmatrix} 1 & a & a & a & \cdots & a \\ -1 & x-a & 0 & 0 & \cdots & 0 \\ -1 & 0 & x-a & 0 & \cdots & 0 \\ -1 & 0 & 0 & x-a & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ -1 & 0 & 0 & 0 & \cdots & x-a \end{vmatrix}$$

化为例 3 的形式.

例 5(范德蒙 Vander Monde 行列式)

$$V_n = \begin{vmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^{n-2} & x_1^{n-1} \\ 1 & x_2 & x_2^2 & \cdots & x_2^{n-2} & x_2^{n-1} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & x_{n-1} & x_{n-1}^2 & \cdots & x_{n-1}^{n-2} & x_{n-1}^{n-1} \\ 1 & x_n & x_n^2 & \cdots & x_n^{n-2} & x_n^{n-1} \end{vmatrix}$$

解 行消去法. 第 n-1 列乘以 $-x_n$ 加到第 n 列上, 第 n-2 列乘以 $-x_n$ 加到第 n-1 列上, 一直下去, 1 列乘以 $-x_n$ 加到第 2 列上, 得

$$V_{n} = \begin{vmatrix} 1 & x_{1} - x_{n} & x_{1}^{2} - x_{1}x_{n} & \cdots & x_{1}^{n-2} - x_{1}^{n-3}x_{n} & x_{1}^{n-1} - x_{1}^{n-2}x_{n} \\ 1 & x_{2} - x_{n} & x_{2}^{2} - x_{2}x_{n} & \cdots & x_{2}^{n-2} - x_{2}^{n-3}x_{n} & x_{2}^{n-1} - x_{2}^{n-2}x_{n} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & x_{n-1} - x_{n} & x_{n-1}^{2} - x_{n-1}x_{n} & \cdots & x_{n-1}^{n-2} - x_{n-1}^{n-3}x_{n} & x_{n-1}^{n-1} - x_{n-1}^{n-2}x_{n} \\ 1 & x_{n} - x_{n} & x_{n}^{2} - x_{n}^{2} & \cdots & x_{n}^{n-2} - x_{n}^{n-2} & x_{n}^{n-1} - x_{n}^{n-1} \end{vmatrix}$$

$$= (-1)^{n+1} \begin{vmatrix} x_1 - x_n & x_1^2 - x_1 x_n & \cdots & x_1^{n-2} - x_1^{n-3} x_n & x_1^{n-1} - x_1^{n-2} x_n \\ x_2 - x_n & x_2^2 - x_2 x_n & \cdots & x_2^{n-2} - x_2^{n-3} x_n & x_2^{n-1} - x_2^{n-2} x_n \\ \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ x_{n-1} - x_n & x_{n-1}^2 - x_{n-1} x_n & \cdots & x_{n-1}^{n-2} - x_{n-1}^{n-3} x_n & x_{n-1}^{n-1} - x_{n-1}^{n-2} x_n \end{vmatrix}$$

$$= (-1)^{n-1}(x_1 - x_n)(x_2 - x_n) \cdots (x_{n-1} - x_n) \begin{vmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^{n-2} \\ 1 & x_2 & x_2^2 & \cdots & x_2^{n-2} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & x_{n-1} & x_{n-1}^2 & \cdots & x_{n-1}^{n-2} \\ 1 & x_n & x_n^2 & \cdots & x_n^{n-2} \end{vmatrix}$$

$$= (x_n - x_1)(x_n - x_2) \cdots (x_n - x_{n-1})V_{n-1}.$$

得递推公式
$$V_n = \prod_{i=1}^{n-1} (x_n - x_i) V_{n-1}$$
. 所以 $V_n = \prod_{1 \le j < i = \le n} (x_i - x_j)$.

注 数学归纳法类型 I: $D_n = \alpha D_{n-1} + \beta$.

例 6 $y \neq z$

$$D_{n} = \begin{vmatrix} x & y & \cdots & y \\ z & x & \cdots & y \\ \vdots & \vdots & \ddots & \ddots \\ z & z & \cdots & y \\ z & z & \cdots & x \end{vmatrix} = \begin{vmatrix} x & y & \cdots & y \\ z & x & \cdots & y \\ \vdots & \vdots & \ddots & \ddots \\ z & z & \cdots & y \\ z & z & \cdots & y \end{vmatrix} + \begin{vmatrix} x & y & \cdots & 0 \\ z & x & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots \\ z & z & \cdots & 0 \\ z & z & \cdots & x - y \end{vmatrix}$$
$$= (x - y)D_{n-1} + y(x - z)^{n-1}$$

由于 y, z 有对称性, 故 $D_n = (x-z)D_{n-1} + z(x-y)^{n-1}$, 联立方程可得结果.

注 数学归纳法类型 II:
$$\begin{cases} D_n = aD_{n-1} + b \\ D_n = cD_{n-1} + d \end{cases} (a \neq c)$$

例 7

$$D_n = \begin{vmatrix} a+b & ab & \cdots & 0 & 0 \\ 1 & a+b & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & a+b & ab \\ 0 & 0 & \cdots & 1 & a+b \end{vmatrix}$$

解 按第 1 列展开,可得 $D_n = (a+b)D_{n-1} - abD_{n-2}$.

注 数学归纳法类型 III: $D_n=pD_{n-1}+qD_{n-2}$, 设 a,b 是 $x^2-px-q=0$ 的根, p=a+b,q=-ab. $\begin{cases} D_n-aD_{n-1}=b(D_{n-1}-aD_{n-2})\\ D_n-bD_{n-1}=a(D_{n-1}-bD_{n-2}) \end{cases}$, 化为类型 I, 类型 II.

练习 P_{33} 1 (1), 3(1), 4(2), 5; P_{37-38} 5, 7; P_{46-47} 14, 17, 18 选做 P_{45} , 2, 15, 19